


CPR



Across

- 4 What does CPR stand for?
- 5 If a person is mildly choking they can make sounds and/or ____ loudly.
- 6 After each compression, let the ____ return to it's normal position.
- 7 give up to 5 _____ with the heel of your hand between the infant's shoulder blades.
- 11 If no one has come to help you and you don't have a phone nearby, after 5 sets of 30 _____ and 2 breaths, leave the child to phone 9-1-1.
- 13 When giving breaths, remember to pinch the _____ closed.
- 16 Early recognition and ____ are crucial for survival from cardiac arrest.
- 20 Opening the airway correctly will help keep the _____ out of the way.
- 24 When providing compressions to an infant, put 2 _____ of one hand on the center of the chest, on the breastbone, just below the nipple line.
- 27 Someone who is _____ is one who moves, speaks, blinks, or otherwise reacts when you tap him or ask if he's OK.
- 29 If the object does not come out after r back slaps, turn the infant onto his back, supporting the head and give up to 5 _____ thrusts using 2 fingers.
- 30 Step two from question 7 is tap and _____.
- 32 Step four from question 7 is to phone 9-1-1 and get an _____.
- 33 To check if an infant is responsive, tap the infant on the _____.
- 34 If a shock is needed, ensure no one is touching the person and push the _____ button.
- 36 When an emergency occurs there is a sequence of steps you should follow to determine whether an adult needs CPR. Step one is to make sure the scene is _____.
- 37 Step five from question 7 is to check for _____.
- 40 Put the thumb side of your fist slightly above the _____ button and well below the breastbone, and wrap your arms around the waist so that your hands are in front.
- 41 Step three from question 7 is to shout for _____.
- 42 Repeat giving 5 back slaps and 5 chest thursts until the infant can breathe, cough, or cry, or until the infant becomes _____.
- 44 To open the airway, put one hand on the forehead, and put the fingers of your other hand on the bony part of the _____.
- 45 An _____ is anyone who has reached or has already gone through puberty.

Down

- 1 What does AED stand for?
- 2 Scan the chest for _____ to ten seconds.
- 3 Some words may be used more than once: word bank, fingers, chin, unresponsive, nose, shout, back slaps, chest, compressions, child, sudden cardiac arrest, seven, safe, tongue, infant, foot, touch, shock, chest, five, adult, heart, mouth, two, 120, cardiopulmonary resuscitation, hand, help, CPR, breastbone, abdomen, belly, responsive, thirty, cough, choking, breathing, two, AED, eight, automated external defibrillator
- 8 When push hard and fast in the center of a persons chest, you pump blood to the brain and _____.
- 9 Step six from question 7 is to start _____.
- 10 Push down hard, at least _____ inches.
- 12 Results from an abnormal heart rhythm, causing the heart to quiver, so it can no longer pump blood to the brain, lungs, and other organs.
- 14 To help an infant with a severe airway block first, hold the infant facedown on your forearm, supporting the infant's head and jaw with your _____.
- 15 When you put the AED pads on the chest, make sure they don't _____ each other.
- 17 As soon as an AED gives a shock, immediately resume _____.
- 18 If someone is pregnant or very large, put your arms under the person's armpits and your hands on the lower half of the _____.
- 19 Use adult pads on anyone _____ years or older.
- 21 If your mouth is too small to cover the infants nose and mouth, cover close the mouth and give breaths through the infant's _____.
- 22 When providing breaths to an infant, cover the nose and mouth of the infant with your _____.
- 23 The choking is _____ if the child cannot breathe, if the child has a cough that makes no sound, or if the child can't talk or make any sound
- 25 An _____ is any child under 1 year old.
- 26 For a larger or pregnant person, pull straight back to give _____ thrusts instead of abdominal thrusts.
- 28 Push at the rate of 100 to _____ compressions per minute.
- 31 Give _____ breaths in ten seconds to minimize interruptions in the compressions.
- 35 Give _____ compressions and two breaths.
- 38 Give quick, upward thrusts into the _____.
- 39 If you think someone is choking, ask, "Are you _____? Can I help you?"
- 43 A _____ is someone older than 1 year who has not yet reached puberty.

Word Bank

- | | | | |
|----------------------------------|--------------|-----------------------|-------------------------------|
| mouth | adult | thirty | Cardiopulmonary resuscitation |
| back slaps | nose | compressions | chest |
| 120 | two | five | unresponsive |
| Automated external defibrillator | compressions | word bank | breathing |
| safe | severe | sudden cardiac arrest | nose |
| chest | compressions | belly | touch |
| help | tongue | choking | AED |
| foot | hand | chest | chin |
| Responsive | Breastbone | CPR | two |
| fingers | child | infant | heart |
| eight | cough | abdomen | shock |
| shout | | | |